

Dissociation of explicit and implicit timing in repetitive tapping and drawing movements

HOWARD N. ZELAZNIK¹, REBECCA M. SPENCER¹, AND RICHARD B. IVRY²

¹*Department of Health, Kinesiology and Leisure Studies, Purdue Interdisciplinary Program in Neuroscience*

²*Department of Psychology, University of California at Berkeley*

Four experiments explore the hypothesis that temporal processes may be represented and controlled explicitly or implicitly. Tasks hypothesized to require explicit timing were duration discrimination, tapping, and intermittent circle drawing. In contrast, it was hypothesized that timing control during continuous circle drawing does not rely on an explicit temporal representation, but rather, temporal control is an emergent property of other control processes, i.e., timing is controlled implicitly. Temporal consistency on the tapping and intermittent drawing tasks was related, and performance on both of these tasks was correlated with temporal acuity on an auditory duration discrimination task. However, timing variability of these three tasks was not correlated with timing variability of continuous circle drawing task. These results support the hypothesized distinction between explicit and implicit temporal representations.

Many tasks have clear temporal goals or require precise temporal information in order for successful achievement of the goal of the task. The distance runner establishes target times for each lap prior to the start of a race. The sprinter must anticipate the firing of the starter's gun to get a jump on the competition. How does an individual translate such temporal goals into an action? We will present evidence to indicate that performers, depending upon the nature of the task, can achieve this goal via two processes. One process depends on an explicit representation of the passage of time, what we will call explicit timing. For the other process, the temporal properties of the action may be emergent (Turvey, 1977), in the sense that an explicit representation of time does not directly guide performance (Ornstein, 1969; Zeiler, 1998). We call this timing implicit as a contrast to the explicit timing process. In the examples from track and field, we expect that the sprinter relies on explicit timing to anticipate the start of the race; the distance runner adopts a pace that will produce the target lap times in an emergent manner. Four experiments are presented that lead us to conclude that the timing of movement initiation is an explicit process while the timing of movement duration is an implicit one. First we turn our attention to notions of explicit timing.

One approach to the study of timing involves describing how different neural structures and architectures can "represent" time. For example, Braitenberg and colleagues have proposed a model in which waves of activity across the parallel fibers of the cerebellar cortex provide a range of temporal delays that support coordinated interactions between different limb segments (see Braitenberg, Heck, & Sultan, 1997). Similarly, Rosenbaum's (1998) broadcast theory of timing is based on the idea that the distance traversed across neuron-like units forms a range of temporally-tuned elements.

On a more psychological level, models of internal timing have considered both oscillatory (Treisman, Faulkner, & Naish, 1992) and interval forms of representation (Ivry & Hazeltine, 1995). These models are behavioral, seeking to describe and explain a wide range of temporal phenomena that can be observed in many different tasks. One such task is repetitive tapping. Wing and Kristofferson (1973) proposed a two-process model to account for the variability of the inter-tap intervals. A central timing process is presumed to regulate movement initiation and a motor implementation process is presumed to determine when the actual response occurs. These two processes are hypothesized to contribute independently to overall timing variability, an assumption supported by a large number of studies (see Wing, 1980; Wing, in press). Within the framework of this two-process model, the central mechanism is believed to provide an explicit representation of the target interval.

Ivry, Keele and Diener (1988) provided neurophysiological evidence that central timing and implementation timing processes are independent. Patients with a lesion only in the medial region of a cerebellar hemisphere exhibited increases in implementation timing variance for the ipsilesional hand, compared to the contralesional hand, whereas those patients with localized lateral damage to a cerebellar hemisphere exhibited increases in central clock-like variability, only for the ipsilesional hand compared to the contralesional hand. Moreover, patients with cerebellar lesions also were impaired on a variety of perceptual temporal processing tasks, suggesting that this structure operates as an internal timing system that is exploited for both motor and non-motor tasks that require the precise representation of temporal information (reviewed in Ivry, 1997).

A different approach for exploring whether a common system underlies motor and perceptual timing involves examining the function relating timing variance to the interval to be timed. Across a range of tasks, temporal variability exhibits a scalar property (Gibbon, 1977; Gibbon, Malapani, Dale, & Gallistel, 1997). Variability is proportional to the interval being timed, i.e., a form of Weber's law in the temporal domain. The Weber fraction, that is, the slope of the variability function, is

Thanks to Tricia Martinic for help in data collection in Experiment 1 and thanks to Sarah Crain, Jennifer Marty, and Rethy Subramanian for help in data collection in Experiment 2. Special thanks to Kristi Hall, Beth Carroll, Dawn Chambers, Marie Phillips, and Mary Kaye Rueth for their assistance in data collection for Experiment 4. H. Zelaznik and R. Spencer were supported on grant DC 02527 (NIH), and R. Ivry was supported on NS 30256 and 1778 (NIH) and NSF ECS-9873474. All correspondence should be directed to HNZ. His email address is hzelaz@purdue.edu.

hypothesized to represent the noise properties of the duration-dependent, central component of timing. Ivry and Hazeltine (1995) observed that the variance of auditory duration discrimination and the variance of timing during finger tapping exhibited the same slope. Similarly, Green, Ivry and Woodruff-Pak (1999) reported that the variance in the onset of the conditioned eyeblink response was related linearly to the duration of the interval between the conditioned and unconditioned stimuli, and that the Weber fraction for this function was statistically equivalent to that found for finger tapping. Ivry and colleagues concluded that equivalent Weber relations are the result of the operation of a common representational timing process, what we call explicit timing.

Individual difference studies provide a third approach for exploring the viability of a general internal clock construct. If two tasks share a common timing process, variability in timing on one of those tasks will be predictive of variability in timing on the other task. In other words, there should be a significant positive correlation for timing variability across the two tasks. Individual differences in timing variability are correlated across movements involving different effectors such as finger and arm (Keele & Hawkins, 1982) or arm and jaw (Franz, Zelaznik, & Smith, 1992). Moreover, variability in timing in finger tapping is correlated with duration-discrimination performance (Keele, Pokorny, Corcos, & Ivry, 1985).

The studies described above provide support for the hypothesis that many tasks share a common timing process. However, the motor demands on these tasks appear to be relatively minor. For example finger tapping, arm tapping, jaw movements, and the eyeblink response all involve movement along a single axis of rotation. Furthermore, the individual can satisfy the task demands by controlling the timing of movement initiation. During eyeblink conditioning, the animal must associate the conditioned stimulus and unconditioned stimulus, and learn when to initiate the eyeblink. Similarly, in the repetitive tapping tasks, the subject could adopt a strategy of producing a fixed duration movement, and vary the interval of time between initiations to meet the demands of the task.

It seems reasonable that in such tasks, an explicit representation of time would be important. However, certain tasks might not require an explicit central representation of time to control periodic behavior (Ivry & Hazeltine, 1992; Semjen, 1992). To examine this issue, we tested subjects on two tasks, a finger tapping task and a continuous circle drawing task (Robertson, Zelaznik, Lantero, Bojczyk, Spencer, Doffin, & Schneidt, 1999). The period of motion of the tapping task was the same as the period of motion for circle drawing. Two important results were obtained. First, individual differences in timing variability in tapping were not significantly correlated with individual differences in timing variability in drawing. Second, when the cycle period was varied, the slope of the variability function for tapping and circle drawing was not the same. Tapping had a much steeper slope than circle drawing. In a follow-up study, Zelaznik, Spencer and Doffin (2000) showed that the differences in temporal variability across the two tasks were not due to differences in the preferred rate of motion. Even when allowed to adopt their preferred rate for each task, timing variability in tapping and circle drawing remained uncorrelated. Thus, we have argued (Robertson et al., 1999; Zelaznik et al.,

2000) that not all timing tasks share the same explicit timing process.

In the present set of experiments we explore the idea that the timing of movement initiation processes involves an explicit representation of time, whereas movements that are continuous may not require an explicit temporal representation. We hypothesize that once initiated, these latter movements no longer involve explicit temporal control. Rather, the temporal regularities are emergent, reflecting the operation of a different control process. We refer to this latter type of timing as implicit. The important issue at hand is whether the use of explicit timing depends upon movements that are relatively discrete, or at least not smooth and continuous.

To test this hypothesis, we developed a hybrid circle-drawing task, intermittent circle drawing. The participant was required to insert a pause between each drawing cycle. A metronome prescribed the duration of the circle and the duration of the pause. The participant was trained to produce the two equal epochs. We assumed that the pause would require explicit timing because of the need to set the timing for movement initiation with each cycle. Based on our assumption that a common timing system is invoked for tasks involving an explicit temporal representation, we expected to observe a significant correlation between measures of temporal precision obtained during the repetitive tapping and the intermittent circle drawing tasks. Alternatively, if the difference between tapping and drawing is due to the complexity of trajectory control, then we would expect to find tapping and intermittent circle drawing task to be uncorrelated, similar to what has been observed with tapping and continuous circle drawing (Robertson et al., 1999).

Experiment 1

In Experiment 1 participants performed the repetitive tapping task, attempting to match a target interval of 1000 ms. The participants also performed the intermittent circle drawing task in which the goal was to draw a circle with a 500 ms movement time, followed by a 500 ms pause. Both tasks were performed using the continuation procedure routinely used for studying timing.

Method

Participants. Twenty five (7 male and 18 female) students aged 18-22 volunteered to be paid participants. All participants, except one, were right handed. All had normal or corrected to normal vision and no known neurological impairments. The Purdue University Committee on the Usage of Human Research Subjects approved recruitment of participants and informed consent procedures.

Tasks. There were two tasks in this study. In both, a computer-controlled metronome produced an alternating high-pitch tone (800 Hz) and a low-pitch tone (400 Hz), of 20 ms in duration. The inter-stimulus interval between successive tones was 500 ms. For the repetitive tapping task, the participants were instructed to tap the index finger of their dominant hand on the table coincident with the high-pitch tone (i.e., a cycle duration of 1000 ms). We chose this duration and metronome characteristics based upon pilot testing and the capabilities of our instrumentation interface. Following a synchronization phase, the

metronome was disengaged. During this continuation phase of each trial, the participants were to continue tapping at the prescribed frequency.

For intermittent circle drawing participants were required to complete one circle during the interval between the low-pitch tone and high-pitch tone (i.e., movement time of 500 ms) and to pause during the 500 ms interval between the high-pitch tone and low-pitch tone. Thus, they alternated every half-second between circle drawing and pausing, using the tones to partition the 1000 ms interval during the synchronization phase. They were instructed to continue in this fashion during the continuation phase.

The target diameter of the circle was 8 cm. The participant was not required to trace a circle, but was guided with 4, 1.5 cm diameter circles positioned at 0, 90, 180 and 270 degrees along the imagined circumference of the 8 cm target circle. In Figure 1 we present a schematic representation of the circle template setup. The task required the subject to pass through these guides during the circle drawing task. The guide at 90 degrees was shaded, and served as the location of the "on-time" marker. This is where the participant's pencil needed to be when the appropriate tone was presented. Because of the relatively large size of the tracing guides relative to the circle diameter ($\text{Log}_2(2D/\text{diameter}) = 3.14$), the spatial demands of the task were low, allowing participants to give priority to the temporal aspects of the task. Instructions emphasized that the main goal was to be on time, and the template was to provide a guide for the approximate size of the circle and the location of the temporal synchrony point.

Apparatus. The response apparatus consisted of a 79 cm high table on which a target sheet was secured. A Mars-Staedtler 2 mm pencil holder was used for the circle drawing task. The bottom portion of the pencil was wrapped tightly with adhesive tape resulting in a 1 cm diameter barrel. Attached below the taped portion of the pencil was an infrared light emitting diode (IRED) from a Watsmart kinematic recording system. For the tapping task, a bandaid was wrapped around the nail of the index finger of the dominant hand. The IRED was secured to the bandaid with Velcro.

An MS-DOS based computer was used to generate the metronome signal. The computer was programmed to produce an alternating sequence of high (800 Hz) and low (400 Hz) tones, 50 ms in duration, with a 500 ms interstimulus interval.

Procedures. After informed consent was obtained and the participant was provided with general instructions, testing began. A trial began with the participant placing their finger (for tapping), or pencil (drawing) on the target circle on the tabletop. The experimenter said "ready - begin", and then the metronome was engaged. The participant entrained their movements to the metronome. When the metronome was disengaged the participant attempted to maintain the same rate during the continuation portion of the trial. After a period of time long enough to produce 35 continuation movements, the trial ended with a series of 5 tones with a 50 ms interval. In the drawing tasks, the paper containing the circle template was replaced following each trial. There was a 25 second intertrial interval.

The standard protocol for correlational studies examining individual differences (e.g., Keele et al., 1985; Robertson et al., 1999) was employed. Tasks were performed in a fixed order. Tapping was performed first followed by intermittent circle drawing. Preliminary testing had revealed that producing equal

pause and movement segment durations during continuation was difficult. Thus, we included a training procedure for both tasks consisting of ten trials of 54 cycles each. On training trial one, all of the movements were performed with the metronome. On each of the subsequent nine training trials, the number of synchronization cycles decreased by four, and the number of continuation cycles increased by four. Thus, at the end of the tenth training trial there were 18 synchronization and 36 continuation cycles. A participant then continued with the test phase. This consisted of 10 trials, each composed of 18 synchronization cycles and 36 continuation cycles.

Data Acquisition. A two-camera Watsmart kinematic recording system sampled the location of the IRED at 256 Hz. The three dimensional data series were reconstructed off-line.

Results and Discussion

Data Analysis and Reduction. We used an interactive graphical routine to determine the cycle duration for tapping, and to parse the intermittent circle drawing record into the movement and pause components. This procedure has been shown to be extremely accurate (plus or minus one sample), and reliable (inter-rater reliability, $r > .98$) in identifying specific target points within a movement cycle (see Robertson et al., 1999). The raw displacement record and the filtered displacement record (25 Hz, low-pass, forward and backward) were displayed on a monitor. A mouse-controlled cross-hair was moved to mark the end of the downward movement of the index finger of tapping or the pause and movement components for intermittent drawing. Based on this procedure, we determined the duration of each cycle and retained the middle 31 cycles of the continuation phase for further analysis. A linear regression analysis was performed on these cycles and the measure of timing variability was based on the residuals from the regression line. This procedure eliminates temporal variability that arises from global changes in performance rate (i.e., tendency to speed up or slow down) and has been used in many studies of repetitive tapping (see Keele et al., 1985; Vorberg & Wing, 1996).¹ Because we were interested in variability in timing under the best of conditions, only the six trials with the lowest variability were included in the analyses reported below (see Peters, 1989 for further logic on this issue).

Descriptive Results. In Figure 2 a typical trajectory for four seconds of continuation performance is presented for both tapping and intermittent drawing. A distinct pause can be seen during both tapping and intermittent drawing. The descriptive data are presented in Table 1. The main dependent variable for temporal variability is the coefficient of variation, computed as the standard deviation in timing divided by the mean interval. On average, the participants performed the tasks successfully, with the mean cycle duration close to the target interval of 1000 ms. The observed coefficient of variation values are in accord with previous reports (e.g., Allan, 1979; Ivry & Hazeltine, 1995). The cycle for circle drawing was partitioned into movement duration and pause duration components. Participants had some difficulty in parsing the intermittent circle into equal pause and movement durations. Thus, while the participants were very successful in sustaining the overall duration, they did not maintain a 1:1 ratio of movement and pause time. The variability of these components was larger (in terms of percentages) than the variability of the total cycle duration. Thus, there was greater inconsistency in maintaining the movement and pause subintervals in comparison to the total

interval. Billon, Semjen, and Stelmach (1996) have argued that this type of result is indicative of a hierarchical representation of the action with the primary temporal goal associated with maintaining the overall period.

Descriptive spatial variables also are included in Table 1. For the finger tapping task, we determined the maximum height of the IRED with respect to the table top. For intermittent circle drawing, the center of the circle was computed on a cycle by cycle basis as the mean x and y coordinates. Then the radius for each sample with respect to that center was calculated. For each cycle, the average radius was calculated and averaged over cycles within a trial. Within a cycle the standard deviation of the radius was computed. This is a measure of the integrity of the circle shape.

	TEMPORAL			SPATIAL		
	MT (ms)	CV (%)	Rel	Disp (cm)	CV (%)	Rel
<u>Tapping</u>	1019.6 (11.7)	6.2 (0.25)	.89	3.3 (0.54)	7.1 (1.2)	.85
<u>I Circle</u>						
total	1030.3 (11.9)	5.5 (1.0)	.86			
move	653.9 (14.2)	7.1 (0.49)	.83	3.9 (0.45)	12.3 (0.98)	.88
pause	376.4 (15.7)	13.2 (0.52)	.83			

Table 1. Descriptive data for Experiment 1. Standard errors are in parentheses. I Circle stands for the intermittent circle drawing task, MT is the average time, Disp is the displacement, CV is the coefficient of variation, and Rel stands for reliability.

A smaller standard deviation indicates that the trajectory is more circular. The standard deviation was converted to a coefficient of variation and then averaged across cycles. As seen in Table 1 the average radius of the circle was 3.9 cm and the coefficient of variation was about 12 percent. The average tapping displacement was slightly greater than 3 cm with a coefficient of variation of about 7 percent.

Correlational Analysis. Separate subject by trial analyses of variance (ANOVA) were conducted on each task separately for the best six trials to compute reliability for each task. Reliability, which varies between zero and one, was computed from the ANOVA by computing the ratio of variance due to subjects, relative to variance due to the trials and the subject by trials interaction terms (see Thomas & Nelson, 1996). As seen in Table 1, reliability values were equal to or greater than .83. These values are similar to our previous individual difference work (Franz, et al., 1992; Robertson et al., 1999; Zelaznik, et al., 2000).

The primary analyses centered on the correlation matrix between the variability scores on the two tasks. These are presented in Table 2. With 25 subjects, an r value of .37 is significant at the .05 level. There was a significant correlation between tapping and intermittent circle drawing ($r = .55$). When tapping variability was compared to the components of intermittent drawing, we found that tapping variability was significantly correlated with the variability of the pause subinterval ($r = .55$), but not the variability of the movement subinterval ($r = .21$). However, the two phases of the intermittent task were correlated ($r = .65$).

	TEMPORAL			SPATIAL	
	total	I Circle move	pause	<u>Tapping</u>	I Circle move
TEMPORAL					
<u>Tapping</u>	.55	.21	.55	.30	.23
<u>I Circle</u>					
total	.78	.67	.11	.22	
move		.65	.09	.38	
pause			.21	-.09	
SPATIAL					
<u>Tapping</u>					.08

Table 2. Correlation values for Experiment 1 based on temporal and spatial coefficient of variation. I Circle is the intermittent circle drawing task. Significant correlations ($p < .05$) are in boldface.

It is interesting to note that, for intermittent circles, the variability in the spatial aspect of the circle was correlated with temporal variability of the movement portion of the circle, but not with the pause. Finger tapping timing variability was not related to finger tapping spatial variability. Finally, spatial variability in tapping was not correlated with spatial variability in circle drawing.

The results of the present experiment reveal a significant correlation between temporal variability on the tapping and intermittent circle drawing tasks. This finding contrasts with the earlier report of Robertson et al. (1999) in which temporal variability in continuous drawing and tapping was not correlated. We believe that the discrepancy reflects a fundamental difference in how people achieve a temporal goal when drawing circles in a continuous or intermittent manner. As outlined above, we hypothesize that a series of periodic, discrete movements requires an explicit temporal representation. We propose that when a pause is inserted in the circle task, the participants must utilize an internal timing process to directly control the required interval between movement cycles and to orchestrate movement initiation. This process is assumed to be similar to that invoked in the repetitive tapping task. The pause, by transforming the cycles of the circle task from continuous to discrete, makes the task more similar to tapping in terms of control processes. We hypothesize that this explicit control of timing is used under such discrete conditions, providing the requisite representation for specific events (e.g., the time of contact in tapping, or the duration of the pause or inter-onset times in the circle drawing task).

Experiment 2

To allow for direct comparisons between discrete and continuous tasks, we included a continuous circle drawing condition in Experiment 2 along with the tapping and intermittent circle tasks. By examining the pattern of correlations between the three tasks, we can better examine the proposed distinction between explicit and implicit timing. Based on Experiment 1 and our previous work (Robertson et al., 1999; Zelaznik et al., 2000), we expected to obtain a positive correlation between the measures of temporal variability on the tapping and intermittent circle drawing tasks. On the other hand, the correlation should be low between tapping and continuous circle drawing.

Of central interest is the relationship between temporal variability on the two circle drawing tasks. One possibility is that these tasks will be positively correlated given shared processes associated with trajectory formation and/or processes involved in

the visual control of circle drawing. However, based on our assumption that the intermittent circle task entails explicit temporal control due to the timing requirements of movement initiation, we predicted that the correlation between the two circle drawing tasks would be low despite their superficial similarity.

A second change in Experiment 2 involved the cycle duration. The rate of movement (1 Hz) in Experiment 1 was slower than typically used in repetitive tapping studies and slower than that used in the experiments of Robertson et al. (1999). It is possible that the observed correlation between discrete circle drawing and tapping in Experiment 1 was due to the use of this slow rate rather than the intermittency of the circle drawing task. For example, perhaps feedback processes become more relevant as the cycle duration increases and the correlational results reflect individual differences in the use of feedback. Thus, we adopted a cycle duration of 800 ms in Experiment 2, which was one of the rates used in the studies of Robertson et al. (1999).

Method

Participants. Twenty five, right-handed college-aged adults (14 males and 11 females) served as paid volunteer research participants. Informed consent was obtained from all participants according to procedures approved by the Purdue University Committee on the Usage of Human Research Subjects.

Apparatus, Tasks, and Procedures. To ensure that the participants could complete the circle in the targeted time, the circle template on the target sheet was reduced to 5 cm in diameter. Only one tracing guide (a small circle, 1 cm in diameter) was shown on the circumference of the 5 cm diameter circle, positioned at 90 degrees (see bottom of Figure 1). The tapping and intermittent circle drawing tasks were the same as in Experiment 1, except for the change in target frequency. The inter-stimulus interval for the high- and low-pitched tones was set to 400 ms. The tapping task and intermittent circle drawing were performed in a manner identical to Experiment 1. For the continuous circle task, the participants were to complete one circle every two beats (800 ms movement time), moving continuously between cycles.

The participants first completed a training phase structured in a similar manner as in Experiment 1. During the test phase, there were ten trials, each consisting of 18 paced cycles and 36 unpaced cycles. A fixed order was used with the tapping condition performed first, then the intermittent circle drawing, and finally continuous circle drawing.

Data Collection. Kinematic three-dimensional displacement data were collected with an Ascension Technology, mini-Birdtm magnetic system, which sampled the location of the markers at 140 Hz.

Results and Discussion

Data Reduction and Analysis. As in Experiment 1, a graphical analysis routine was used to score the tapping and intermittent circle drawing data. Because the continuous circle drawing task was performed very smoothly without pauses, the kinematic data were low pass filtered (8 Hz cutoff, forwards and backwards). The displacement records along the y axis were differentiated and a numerical algorithm was used to determine when the velocity in the y dimension changed sign (a movement reversal). The interval between reversal n , and reversal $n+2$ was recorded as the duration of one complete cycle. Regression analyses were used as

in Experiment 1 to remove global changes in movement rate, and the variability measures were based on the detrended time series. The best six trials from the test phase were included in the analyses.

Descriptive Data. In Figure 3 we present representative kinematic traces for each of the three tasks. Both the tapping and the intermittent circle tasks show discernable pauses. The pauses during tapping were uninstructed, reflecting the natural mode that participants adopt when tapping at the 800 ms rate; for the intermittent drawing task, the instructions specified that a pause should be inserted between each movement cycle. In contrast, the continuous circle drawing trajectory is very smooth, without any discernable pauses.

The descriptive data are presented in Table 3. The participants were able to meet the demands of each task. In contrast to Experiment 1, the participants were able to parse the full cycle into pause and movement phases that were of approximately equal duration in the intermittent circle drawing task. For all three tasks, reliability values were high and the coefficient of variation

	TEMPORAL			SPATIAL		
	MT (ms)	CV (%)	Rel	Disp (cm)	CV (%)	Rel
<u>Tapping</u>	803.5 (5.45)	5.9 (0.27)	.92	3.34 (0.66)	10.1 (1.0)	.89
<u>C Circle</u>	784.9 (8.98)	5.9 (0.14)	.90	2.54 (0.20)	6.3 (0.36)	.91
<u>I Circle</u>						
total	823.4 (8.12)	5.7 (0.21)	.91			
move	420.4 (5.55)	8.6 (0.23)	.86	2.62 (0.24)	17.9 (1.6)	.88
pause	403.1 (7.45)	10.7 (0.39)	.83			

Table 3. Descriptive data for Experiment 2. Standard errors are in parentheses. I Circle stands for the intermittent circle drawing task, MT is the average duration, Disp is for displacement, CV is the coefficient of variation, and Rel stands for reliability.

on the temporal measure was less than 6 percent. For the spatial domain, the circles were drawn about the appropriate size for both drawing tasks. However, the measure of spatial variability was much lower for the continuous circles compared to the intermittent circles.

Correlational Analysis. In Table 4 we present the correlation values based on the coefficient of variation measures (critical r value is .37, $p < .05$). Replicating Experiment 1, temporal variability in tapping and intermittent circle drawing tasks was significantly correlated ($r = .50$). However, while positive correlations were obtained between continuous circle drawing and the tapping and intermittent circle tasks, neither value was significant. Thus, individual differences in temporal control were significantly related between the two tasks hypothesized to require explicit temporal control. In contrast, the correlations were reduced between these tasks and the continuous drawing task.

In Table 4 the correlation matrix based on temporal variability associated with the movement and pause phases of the intermittent drawing task, also are presented. Total variability on this task is more highly correlated with the pause phase than the

movement phase. This result is consistent with the hypothesis that explicit temporal control is related to timing the onset of each cycle, and it is the pause phase in which such control is most required. Further support comes from the fact that the pause phase during intermittent circle drawing was correlated with tapping variability whereas the movement phase was not. However, contrary to our expectations, the pause phase was also correlated with variability on the continuous drawing task, although to a lesser degree than with tapping ($r = .39$ vs. $.53$). The overall pattern of positive correlations observed in this experiment suggest that there is some shared timing factor associated with all three tasks (and utilized to a lesser extent in continuous circle drawing), or that part of the individual difference variation is related to general factors.

The correlation values for the spatial and temporal measures reveal some interesting patterns. In general these correlations are positive, yet low, indicating that factors contributing to temporal and spatial variability are to some extent dissociable. However, it is interesting to note that the few values that approached significance involved the correlation between measures of spatial and temporal variability on the same circle drawing tasks. In

	TEMPORAL			Tap	SPATIAL	
	C Circle	I Circle total move	pause		C Circle	I Circle move
TEMPORAL						
Tap	.28	.50	.18	.53	.10	.18
C Circle		.27	.32	.39	-.01	.33
I Circle						
total			.47	.81	.09	.22
move				.27	.20	.36
pause					-.03	.13
SPATIAL						
Tap						-.18
C Circle						.09
						.39

Table 4. Correlation values for Experiment 2 based on temporal and spatial coefficient of variation. I Circle stands for the intermittent circle drawing task, C Circle stands for the continuous circle drawing task. Significant correlations ($p < .05$) are in boldface.

addition, a significant correlation was found between the measures of spatial variability on the two circle drawing tasks.

The low of correlation, for temporal variability, between the two circle drawing tasks is noteworthy for three reasons. First, at a superficial level, the continuous and intermittent circle drawing tasks appear to be the most similar of the three tasks. Both require multi-joint movements and the task goal entails not only the production of periodic movements, but movements that conform to a specified trajectory. In contrast, tapping involves a single joint movement and there is little reference to a target trajectory. Nonetheless, temporal variability was more closely related between the tapping and intermittent drawing task.

Second, the results fail to support the hypothesis that the low correlations between tapping and continuous drawing tasks, both in the current experiments and in Robertson et al. (1999), arise because variability on the drawing tasks reflects noise in processes associated with trajectory planning and execution. One might suppose that such factors are relevant for the circle drawing tasks. The circle requires more degrees of freedom (greater number of involved joints) than tapping and the use of on-line visual guidance is likely to be higher. If these factors were

relevant, we would expect the continuous circle drawing task to be more highly correlated with intermittent circle drawing than tapping. This was not the case. Continuous circle drawing was modestly correlated with both of the other two tasks, but this non-significant correlation was no higher for intermittent drawing task than for tapping.

Third, the lack of correlation between the continuous drawing task and the intermittent drawing (and the tapping task) suggests that the individual differences underlying the correlation matrix are not associated with a general factor that applies across all tasks. In summary, the major results of Experiment 2 were in accord with the predictions derived from the explicit-implicit timing distinction. The two tasks that we hypothesized required explicit timing, those involving non-continuous responses, exhibited significant correlations on the measure of temporal variability. In contrast, the correlations were reduced, and failed to reach significance between these two tasks and the continuous circle drawing task. Taken together, Experiments 1 and 2 demonstrate the importance of movement continuity in accounting for individual differences in temporal variability on repetitive movement tasks. We believe that this factor is critical for the distinction between explicit and implicit temporal control in the current set of tasks. We hypothesize that the timing of successive cycle onsets forms the basis for an explicit temporal representation when the task involves a series of relatively discrete responses. In contrast, timing may become secondary to other control parameters that are used to ensure that the movements remain continuous and smooth.

Experiment 3

The hypothesis that tapping and discrete circle drawing draw on a common explicit timing process was supported in the first two experiments. We have proposed that in such movements, the temporal goal is explicitly represented, providing a signal for when each relatively discrete movement should be initiated. In Experiment 3 we turn to the relationship of these tasks with a non-motor task in which explicit timing is required.

Keele et al. (1985) found that performance on an auditory duration discrimination task was correlated with timing precision in tapping ($r = .50$). Keele et al. inferred that these two tasks share a common timing mechanism. We replicate and extend this work in Experiment 3, examining the correlation matrix between those two tasks as well as our continuous and intermittent circle drawing tasks. We predict that tapping, intermittent circle drawing, and duration discrimination will all be positively correlated, reflecting the fact that each draws upon an internal timing system required for providing explicit temporal representations. In contrast, we predict that the correlations will be lower or even absent between variability on the continuous circle drawing and the other three tasks. This prediction provides a strong test of the hypothesized distinction between tasks involving explicit and implicit timing given the substantial differences between the tapping, intermittent drawing, and duration discrimination tasks. As such, Experiment 3 provides an important generalization of the extent of explicit timing in motor and non-motor tasks.

Method

Participants. Thirty-five students (21 female, 14 male) aged 18-24 volunteered for the experiment in exchange for payment. Informed consent was obtained from all participants according to procedures approved by the Purdue University Committee on the Usage of Human Research Subjects.

Apparatus. The auditory duration discrimination task was run on an MS-DOS based computer. An amplifier and speaker controlled by the computer were positioned behind the subject. The apparatus for the temporal production tasks was the same as those used in Experiment 2.

Tasks and Procedure. All of the temporal production tasks were administered as in Experiment 2. The duration discrimination task was identical to that used by Keele et al. (1985). In brief, each trial contained four tones, grouped into two pairs of two tones with the pairs separated by 1 s. The first pair formed the standard interval and was always separated by 400 ms. The interval between the second pair varied, constituting the comparison interval. Participants judged whether the comparison interval was shorter or longer than the standard interval by pressing the "S" key on the computer keyboard if the comparison interval was perceived as shorter and the "L" key if the comparison interval was perceived as longer than the first.

The comparison interval was set to a value that was either shorter or longer than the standard interval based on an adaptive parameter estimation by sequential testing (PEST) procedure (Ivry & Hazeltine, 1995). The PEST procedure sets the comparison value to the current estimate of the participant's difference threshold. In the current experiment, the estimated threshold corresponded to one standard deviation on the assumed psychophysical function, or a point at which participants would be correct on approximately 72% of the trials. Separate estimates were made for the threshold in which the comparison interval was shorter than the standard and in which the comparison was longer than the standard, although the trials used to estimate these thresholds were intermixed within the test block. The psychophysical function was recalculated after each response, providing a new estimate of the difference threshold and thus generating the test value for the next trial. The threshold estimates after 60 trials (30 per threshold) provided a measure of temporal acuity. The difference between the two estimates corresponds to a range of two standard deviations on the psychophysical function and we use half this value (1 SD) as our measure for the analyses. The test trials were preceded by 11 practice trials. The first ten of these involved a set of fixed durations, selected to be easily discriminable. The comparison interval for the last practice trial was equal to the standard interval (400 ms) to minimize between-trial bias.

The task order was fixed. The order was duration discrimination for one trial block, tapping, continuous circle drawing, intermittent circle drawing and a second block of duration discrimination. The entire experiment lasted approximately 85 min.

Data Collection. A miniBIRD™ kinematic data collection system sampled the three dimensional location of the magnetic markers at 140 Hz. Only kinematic data from the continuation portion of the testing trial was collected.

Results and Discussion

Descriptive Data. One participant with scores clearly outside the range of the other subjects was not included in the analysis. This participant's average cycle duration was 557 ms for the movement portion of intermittent circles, 100 ms greater than that of any other participant. Analyses of the data from the remaining 34 participants were based on the best six testing trials in each movement task as determined by the detrended variance. The lower part of Figure 3 shows a sample of representative tapping data (at 400 ms rate) for this experiment. Even at this more rapid rate of tapping there are discernable pauses in the tapping movements.²

In Table 5 the movement time and coefficient of variation for all of the temporal production tasks are displayed. Participants produced appropriate average durations, with the exception of intermittent circles, which varied slightly from the goal time of 400-ms circle intervals followed by a 400 ms pause intervals. The mean coefficients of variation were in the 5-6 percent range, consistent with previous studies and Experiments 1 and 2. Moreover, the coefficient of variation for the pause and movement of the intermittent circle task was greater than the coefficient of variation values for the total cycle time. The radius for intermittent circles was about the same value as that of continuous circles, but as found in Experiment 2 the intermittent circles had about 50 percent more spatial variability than the continuous circles. All reliabilities for the temporal production tasks were greater than .83.

The mean standard deviation of the duration task across the two trial blocks was 36 ms. The point of subjective equality, the value at which participants are equally likely to judge a comparison interval as shorter or longer than the standard, was 415 ms indicating a bias to report the comparison interval as

	TEMPORAL			SPATIAL		
	MT (ms)	CV (%)	Rel	Disp (.cm)	CV (%)	Rel
<u>Tapping</u>	404.7 (2.8)	6.6 (0.30)	.94	2.9 (0.26)	13.3 (0.99)	.90
<u>C Circle</u>	402.9 (6.8)	5.1 (0.27)	.87	2.6 (0.18)	11.1 (0.96)	.97
<u>I Circle</u>						
total	862.3 (11.3)	5.3 (0.28)	.90			
move	419.1 (8.9)	6.4 (0.35)	.90	2.4 (0.30)	15.6 (2.1)	.85
pause	441.4 (9.8)	9.5 (0.50)	.83			

Table 5. Descriptive data for Experiment 3. Standard errors are in parentheses. I Circle and C Circle stands for the intermittent and continuous circle drawing task, respectively. MT is the average duration, Disp is for displacement, CV is the coefficient of variation, and Rel stands for reliability.

shorter than the standard. Our focus in the correlational analyses below is restricted to the standard deviation scores because these provide an estimate of temporal variability. As with the production tasks, reliability was quite high on the discrimination task (.79).

Correlation Analysis. In Table 6 the correlation matrix for the various measures from the four tasks is presented. With 34 participants, correlation scores greater than .34 are significant at the .05 level.

	TEMPORAL				SPATIAL			
	<u>C Circle</u>	<u>I Circle</u>	<u>Percept</u>	<u>Tapping</u>	<u>C Circle</u>	<u>I Circle</u>	<u>Percept</u>	<u>Tapping</u>
	total	move	pause		total	move	pause	
TEMPORAL								
<u>Tapping</u>	.26	.40	.52	.39	.39	.22	-.09	.14
<u>C Circle</u>		.18	.30	.16	-.15	.11	.40	.21
<u>I Circle</u>			.84	.63	.56	.18	.11	-.01
total				.35	.29	.12	.33	.42
move					.55	.24	.06	-.09
pause								
SPATIAL								
<u>Discrimination</u>						.13	.13	.08
<u>Tapping</u>							.36	.12
<u>C Circle</u>								.61

Table 6. Correlation values for Experiment 3 based on temporal and spatial coefficient of variation. I Circle and C Circle stands for intermittent and continuous circles, respectively. Percept stands for the perceptual timing discrimination task. Significant correlations ($p < .05$) are in boldface.

Consider first the correlations among the temporal measures. As can be seen in the top row, timing variability on the tapping task was significantly correlated with timing variability on both the intermittent circle drawing task and the duration discrimination task. Performance during tapping was also positively correlated with continuous circle drawing, but the value was lower and not statistically significant. In contrast, the correlations between the continuous circle drawing task (the second row) and the various measures of variability on the intermittent drawing task were all non-significant. Most interesting, while no correlation was found between the continuous drawing task and the discrimination task ($r = -.15$), the correlation for total variability on the intermittent drawing task and the discrimination task was high ($r = .56$). Finally, the correlation between the duration discrimination and temporal variability of the pause phase was $.55$.

The correlations between the spatial and temporal variability. In summary, the correlation matrix indicates that individual differences in temporal variability on the continuous circle drawing task are not related to factors associated with the other three tasks. Performance on the continuous drawing task failed to correlate with measures of temporal variability on tapping, intermittent circle drawing, and duration discrimination. In contrast, the latter three tasks were all significantly correlated with each other.

This pattern of results provides support for two hypotheses. First, the control of the cycle duration in continuous drawing is different from that of intermittent drawing. Second, tapping, intermittent drawing, and duration discrimination tasks draw upon a common timing process, and this process is distinct from that used to control timing in the continuous circle drawing task. We believe that the commonality among tapping, intermittent drawing, and duration discrimination reflects the fact that all three require an explicit representation of temporal information. The observed correlations result from individual differences in the noise associated with this internal timing system.

Experiment 4

The significant correlations between temporal variability in tapping and intermittent circle drawing in Experiments 1-3 were

measures were very similar to those found in the previous experiments. Spatial and temporal variability were related for continuous circles. For the intermittent drawing task, these measures were only correlated when considering the movement phase ($r = .42$) and not for either total temporal variability or the pause duration variability. Spatial variability in continuous circles was positively related ($r = .33$) to temporal variability of the movement segment of intermittent circles, although this value was only marginally significant. Spatial variability of intermittent circles was related to spatial variability of continuous circles ($r = .61$). The corresponding value in Experiment 2 was $.39$, suggesting a shared process associated with trajectory control. Surprisingly, spatial variability on the tapping and continuous circle drawing task was significantly correlated in this experiment ($r = .36$); in Experiment 2 this correlation was $-.18$. Given the lack of consistency across experiments, we assume this result is spurious.

observed over a range of target intervals. Tapping was tested at 1000 ms, 800 ms, and 400 ms. The cycle duration of the intermittent task has been either 1000 ms or 800 ms, with target movement phases of either 500 ms or 400 ms, respectively. The consistent pattern of results indicates that the correlations are not dependent on specific intervals, at least within the range explored here. Moreover, the continuous circle drawing task has been performed with target intervals ranging from 400 to 800 ms and at all rates, failed to produce reliable correlations with the other two production tasks.

In our final experiment we explore whether there is any specificity in the individual differences in the duration discrimination task relative to the intermittent circle drawing task and tapping. Participants were tested on two versions of the duration discrimination task, one in which the standard interval was 400 ms and a second in which the standard interval was 800 ms. In addition, the participants performed the tapping task with an 800 ms target interval and an intermittent drawing task with an overall duration of 800 ms, partitioned into a 400 ms pause and 400 ms movement phase. This design allowed us to examine whether the correlation pattern varied as a function of the intervals being used for the three tasks. Specifically, will the correlations between duration discrimination and intermittent circle drawing be highest when the interval for the two tasks

matches in terms of overall duration, or the duration of the intermittent movement's components? The inclusion of a second duration discrimination task also allowed us to replicate the findings of Experiment 3 with a second duration. To limit the number of tasks, we did not include tapping at 400 ms and the continuous circle drawing task.

Method

Participants. Thirty graduate and undergraduate students (24 female, and six male) volunteered to be paid participants in this experiment. Informed consent was obtained from all participants according to procedures approved by the Purdue University Committee on the Usage of Human Research Subjects.

Tasks and Procedure. The procedures were the same as in Experiment 3. The duration discrimination task was performed with standard intervals of 400 ms and 800 ms. For both temporal production tasks, the metronome period was set to 400 ms. Subjects were instructed to tap coincident with every other beat in the tapping task and to complete the movement phase between two successive tones in the intermittent drawing task. For each task there were 15 trials, five training and 10 test trials. For the first trial the metronome was engaged for all cycles. Trials 2 through 5 were paced for 45, 36, 27, 18 movement cycles (90, 72, 54, and 36 metronome beats) and were followed by a sufficient period of time to allow 9, 18, 27, and 36 movement cycles, respectively. On test trials, the metronome was engaged for the first 18 cycles (36 beats) followed by an unpaced phase of 28.8 s (equal to 72 beats). Subjects produced approximately 36 additional movement cycles during this phase.

The tasks were performed in a fixed order as follows: 800-ms-discrimination (trial 1), 400-ms-discrimination (trial 1), finger tapping, intermittent circle drawing, 800-ms-discrimination (trial 2), 400-ms-discrimination (trial 2).

Subjects were given a 25 s inter-trial rest interval and a 5-min rest interval between tasks. During this period between tasks, detailed instructions for the following task were administered. The experiment lasted approximately 70 min.

	TEMPORAL			SPATIAL		
	MT (ms)	CV (%)	Rel	Disp (cm)	CV (%)	Rel
<u>Tapping</u>	826.5 (8.0)	7.1 (0.30)	.96	1.9 (0.25)	9.8 (0.74)	.78
<u>I Circle</u>						
total	864.3 (8.9)	6.1 (0.28)	.90			
move	501.0 (8.4)	11.0 (0.50)	.84	2.4 (0.03)	18.4 (1.8)	.90
pause	363.4 (7.9)	7.4 (0.35)	.88			

Table 7. Descriptive data for Experiment 4. Standard errors are in parentheses. MT stands for average duration, Disp is for displacement, CV coefficient of variation, and Rel stands for reliability coefficient.

Results and Discussion

Descriptive Data. Descriptive variables and reliability coefficients are presented in Table 7. The mean tapping rate was close to the goal of 800 ms. The mean cycle time for the intermittent circle drawing task was considerably longer than the target time, with subjects extending the movement phase of the

cycle. Performance on the 400 ms discrimination task was similar to Experiment 3. The standard deviation for this condition was 38 ms. For the 800 ms condition, the standard deviation rose to 61 ms. Reliability was .78 and .72 for the 400 and 800 ms task, respectively.

Correlation Analysis. The coefficient of variation for the production tasks and the average standard deviation for the perception tasks were used to calculate the correlation matrix. The value for significance is .36 ($n = 30$, $p < .05$). The correlation values are presented in Table 8.

As in the preceding experiments, temporal precision in finger tapping and intermittent circle drawing was highly correlated ($r = .65$). The pause portion, and not the movement portion, of the intermittent circle task was correlated with tapping in this experiment similar to what was found in Experiments 1 and 2. Variability measures of tapping and overall cycle duration on the intermittent circle tasks were significantly correlated with the two versions of the perception task. Furthermore, performance on the two perception tasks were correlated. Thus, the results provide additional support for the hypothesis that all three tasks utilize a common process, one we have proposed is involved in the explicit representation of temporal information.

Experiment 4 allowed us to further assess whether the correlations varied as a function of the represented interval. Interestingly, two results suggest that the correlations were highest for common intervals across the perception and production tasks. First, 800 ms tapping was more strongly correlated with the 800 ms version of the perception task than the 400 ms version. Second, the reverse pattern was found for the correlations between duration discrimination and the pause phase of the intermittent circle drawing task. Here, the higher correlation was found with the 400 ms version of the perception task. Though limited by the fact that we did not include a 400 ms tapping condition and intermittent circle drawing task with an 800 ms pause, temporal specificity may be an additional factor with explicit timing in understanding these significant correlations.

The only significant correlation between temporal and spatial variability was within the movement phase of the intermittent circle drawing task. Similar correlations also were observed in the first three experiments, indicating some shared process is involved in regulating the spatial and temporal properties of a trajectory. As expected, the spatial measures were not related to the duration discrimination performance.

General Discussion

Based upon the results of the four experiments the hypothesis that there is an important distinction between the control processes associated with timing tasks involving discrete and continuous events has been supported. Temporal consistency on the tapping and intermittent circle drawing tasks was significantly correlated in all of the experiments, and performance on these tasks also was correlated with acuity on a duration discrimination task. In contrast, the correlations were either weaker or absent between these tasks and the continuous circle drawing task. While all of the tasks could be considered "timing" tasks, in a broad sense, there are considerable differences between them: motor versus perceptual tasks, one dimensional versus two-dimensional trajectories, single-joint versus multi-joint movements, distal versus proximal muscles. Despite these

differences, the pattern of correlations is in accord with the hypothesized distinction between tasks in which the temporal representation is explicit and those in which timing is implicit.

Admittedly, the distinction between explicit and implicit timing is not crystal clear. Unlike the vast literature on memory, we do not mean to imply anything about a difference between conscious and unconscious knowledge. Rather, we adopt these terms to refer to the manner in which temporal information is represented. For tasks involving explicit timing, we propose that an explicit representation of the temporal goal is required and is used to produce appropriately timed intervals. In tapping and intermittent circle drawing, the goal is in terms of “when” some salient point is achieved within the movement cycle. This might be the point of movement initiation; alternatively, at least in tapping, it could correspond to the point of contact with the table surface (Billon et al., 1996). For either case, we assume that an explicit temporal representation is used to guide performance during each cycle.

In contrast, we propose that implicit timing during a task such as continuous circle drawing, involves what we call the “how” of timing. We assume that the timing characteristics of these movements are the result of the motor system controlling another kinematic and/or kinetic variable. This idea is derived from the idea that timing is an emergent process (Turvey, 1977). Temporal regularities may emerge as subjects attempt to minimize kinematic (Hogan, 1984) or neural (Kawato, 1996) variables. For example, consider the case of a baseball batter attempting to hit a home run. The batter attempts to generate more kinetic energy by swinging faster. The fact that the movement time of the bat is decreased does not imply that the movement time of the bat is directly controlled. Rather, it is just as likely that the speed of the bat results in a particular movement time. Turvey (1977; see also Zeiler, 1998) has argued that one need not postulate the explicit control of time for events that unfold in a precise temporal manner. We believe that temporal regularities may emerge as a function of the higher-level conceptualization of the task. When drawing ellipses, for example the control parameter may center on the relationship between curvature and velocity (Viviani & Terzuolo, 1980; 1982). Circles, having constant curvature, can be produced by keeping the magnitude of tangential velocity constant. Movement rate in such tasks may be controlled by varying the stiffness of orthogonal oscillators in a continuous manner. Temporal regularities would thus be an emergent property in that cycle duration is not directly represented.

The term explicit timing and implicit timing refer to the underlying processes involved in controlling time, not the requirements of the tasks. For intermittent drawing, continuous drawing and tapping the temporal goal is identical; the goal is to complete a cycle of movement at the prescribed duration and to be as accurate and consistent as possible in producing the temporal requirements. We believe that the participant meets this goal for continuous drawing in a manner in which an explicit representation of an individual temporal interval is not required.

However, an explicit temporal representation is likely part of the overall goal in our continuous circle drawing task. The target interval is externally specified and subjects are quite adept in adjusting their performance to match this target. However, we believe that people transform this explicit temporal goal into a different control variable once the task is initiated. For example, achieving a circling period of 800 ms rather than 1000 ms may

require control and modulation of the stiffness of the limb. There no longer exists a need to maintain an explicit temporal representation once the motor execution processes are put into play. On the other hand, the temporal goal may still persist, perhaps providing a means for monitoring performance, over several cycles to determine if the cycle durations drifted from the desired target duration. This hypothesis may account for the fact that we observed positive, albeit, non-significant correlations between continuous circle drawing and the two explicit movement tasks in Experiments 2 and 3.

The introduction of the intermittent circle drawing task provided an important extension of the findings of Robertson et al. (1999). In that study, temporal precision in tapping and continuous circle drawing was not related even though the period of motion was equivalent for the two tasks. This null result led to the proposal that these tasks use different timing processes. The intermittent circle drawing task was designed to test whether this difference reflected the differential reliance on explicit timing. Intermittent circle drawing is a hybrid of tapping and continuous circle drawing. Like continuous circle drawing, the intermittent task requires the participants to produce a specified trajectory in addition to a constant rhythm. Moreover, if such trajectories involve visual guidance, the demands here would be similar for the continuous and intermittent versions [see Zelaznik & Lantero, 1996 concerning the role of vision in timing circle drawing movements]. However, similar to the natural strategy adopted during tapping, the intermittent task entails a pause between successive cycles.

The correlation between tapping and intermittent circle drawing also indicates that the critical factor is not related to the complexity of the movements. For example, one might hypothesize that movements along one axis of motion may be different than those that require coordination along multiple axes of motion. This idea is not consistent with results showing a modest correlation between line drawing and circle drawing (Robertson et al., 1999). Furthermore, temporal variability in line drawing and tapping were unrelated (Robertson et al., 1999), even though both movements are one dimensional. We obtained significant correlations between tapping and intermittent circle drawing despite the fact that circle drawing involves two-dimensional movements and tapping only one. Thus, spatial complexity does not appear to be a relevant factor.

The current results have important implications for the use of tapping as the “preparation” in the study of timing. We believe tapping is an excellent task to examine explicit timing processes (see Wing, in press), but cannot serve as a general model for timing in motor control even when there is a specific temporal goal. The implicit timing processes which we believe are emergent must be accounted for in continuous drawing tasks. The temporal properties of continuous drawing tasks likely emerge from the interaction between centrally driven components and the local bone-muscle machinery (see Kugler, Turvey, & Kelso, 1982). Under such conditions, a central explicit component of timing might be hidden from direct examination due to non-linearities in the peripheral processes (Heuer, Schmidt, & Ghodsian, 1995).

The results involving the perceptual tasks in Experiments 3 and 4 offer converging evidence for the hypothesized explicit-implicit distinction. Significant correlations were observed between the perception tasks, tapping, and intermittent circle

	TEMPORAL				SPATIAL		
	total	I Circle move	pause	Discrimination 800	400	Tapping	I Circle move
TEMPORAL							
Tapping	.65	.05	.50	.57	.36	.32	.01
I Circle							
total		.41	.76	.47	.56	.24	.29
move			.39	.17	.25	.26	.41
pause				.32	.66	.12	.07
Discrimination							
800					.54	.25	.06
400						.09	.02
SPATIAL							
Tapping							.18

Table 8. Correlation values for Experiment 4 based on temporal and spatial coefficient of variation. I Circle and C Circle stand for the intermittent and continuous drawing tasks, respectively. Significant correlations ($p < .05$) are in boldface.

drawing. In contrast, no significant correlations were found between the perception tasks and continuous circle drawing. These results are important for two reasons. First, the dissociation provided a strong test of predictions derived from the explicit/implicit hypothesis. Second, when conducting individual difference research, it is important to verify that positive correlations are not obtained across all tasks, to be certain that one has not captured a general factor such as motivation across these tasks.

In Experiment 4 the strongest relationship among the tasks involving explicit timing was found between temporal measures based on similar intervals. The 800 ms tapping task exhibited specificity with respect to duration discrimination. The correlation was stronger between tapping and 800 ms discrimination than 400 ms discrimination. Furthermore, we observed a stronger correlation between 400 ms-discrimination and the pause of intermittent drawing (400 ms). One possibility is that explicit timing in the intermittent circle drawing task is restricted to the pause phase, with this interval being summed to that produced by the implicitly timed movement phase. Alternatively, the intermittent task may entail a hierarchical temporal representation with a top level representing the time between successive movement onsets and a subordinate timer used to impose the pause interval. The latter hypothesis is supported by the fact that the coefficient of variation is substantially lower when measured for the total interval compared to the coefficient of variation of the pause phase.

An important question for future research concerns the neurological basis for the distinction between explicit and implicit timing. Ivry (1996; 1997; Ivry & Keele, 1989) has argued that the cerebellum is essential for tasks requiring a precise temporal representation such as repetitive tapping, duration discrimination, and eyeblink conditioning. As such, the cerebellum would be an essential structure involved in explicit timing. It is, however, possible that the cerebellum may also be involved in the control of tasks for which timing may be implicit. For example, patients with cerebellar ataxia are impaired on tasks involving complex trajectories such as throwing (Martin, Keating, Goodkin, Bastian, & Thach, 1996; Timmon, Watts, & Hore, 1999). A detailed analysis of the kinematics and EMG records during these movements indicates that their impairment appears to be in the

fine temporal coordination between the gestures of different limb segments.

What may change on tasks involving implicit timing is the contribution of other processes to overall temporal variability. For example, temporal consistency during continuous circle drawing may reflect processes associated with trajectory formation and control. These processes may be associated with non-cerebellar structures. Indeed, the neurological literature suggests that spatial planning arises at a cortical level (e.g., Franz, Eliassen, Ivry, & Gazzaniga, 1996; Desmurget, Epstein, Turner, Prablanc, Alexander, & Grafton, 1999). Perhaps variability with this aspect of coordination dominates performance during continuous circle drawing.

Footnotes


1. The drift was minor in almost all cases, yielding a detrended variance that was just slightly lower than that obtained from the raw data.
2. That the 400 ms tapping movements show a pause even without having to tap on every other beat as in the 800 and 1000 ms tasks, leads us to believe that the pause is a natural component of the tapping movement and not an artifact of the metronome characteristics of the first two experiments.

References

- Allan, L. G. (1979). The perception of time. *Perception and Psychophysics*, *26*, 340-354.
- Billon, M., Semjen, A., & Stelmach, G. E. (1996). The timing effects of accent production in periodic finger-tapping sequences. *Journal of Motor Behavior*, *28*, 198-210.
- Braitenberg, V., Heck, D., & Sultan, F. (1997). The detection and generation of sequences as a key to cerebellar function: Experiments and theory. *Behavioral & Brain Sciences*, *20*, 229-277.
- Desmurget, M., Epstein, C. M., Turner, R. S., Prablanc, C., Alexander, G. E., & Grafton, S. T. (1999). Role of the posterior parietal cortex in updating reaching movements to a visual target. *Nature Neuroscience*, *2*, 563-567.
- Franz, E. A., Eliassen, J. C., Ivry, R. B., & Gazzaniga, M. S. (1996). Dissociation of spatial and temporal coupling in the bimanual movements of callosotomy patients. *Psychological Science*, *7*, 306-310.

- Zelaznik, Spencer, & Ivry
 Journal of Experiment Psychology: Human perception & Performance (2002), v 28: 575-588
 Timing Processes
- Franz, E. A., Zelaznik, H. N., & Smith, A. (1992). Evidence of common timing processes in the control of manual, orofacial, and speech movement. Journal of Motor Behavior, *24*, 281-287.
- Gibbon, J. (1977). Scalar expectancy theory and Weber's law in animal timing. Psychological Review, *23*, 279-325.
- Gibbon, J., Malapani, C., Dale, C. L., & Gallistel, C. R. (1997). Toward a neurobiology of temporal cognition - Advances and challenges [Review]. Current Opinion in Neurobiology, *7*, 170-184.
- Green, J. T., Ivry, R. B., & Woodruff-Pak, D. S. (1999). Timing in eyeblink classical conditioning and timed-interval tapping. Psychological Science, *10*, 19-23.
- Heuer, H., Schmidt, R. A., & Ghodsian, D. (1995). Generalized motor programs for rapid bimanual tasks: A two-level multiplicative-rate model. Biological Cybernetics, *73*, 343-356.
- Hogan, N. (1984). An organizing principle for a class of voluntary movements. Journal of Neuroscience, *4*, 2745-2754.
- Ivry, R. (1997). Cerebellar Timing Systems. In J. Schmahmann (Ed.), The Cerebellum and Cognition (pp. 555-573). San Diego: Academic Press.
- Ivry, R. B. (1996). The representation of temporal information in perception and motor control. Current Opinion in Neurobiology, *6*, 851-7.
- Ivry, R. B. & Hazeltine, R. E. (1992). Models of timing-with-a-timer. In F. Macar (Ed.), Time, action and cognition: Towards bridging the gap, (pp 183-189). Dordrecht, Netherlands: Kluwer Academic Publishers.
- Ivry, R. B. & Hazeltine, R. E. (1995). Perception and production of temporal intervals across a range of durations: Evidence for a common timing mechanism. Journal of Experimental Psychology: Human Perception and Performance, *21*, 3-18.
- Ivry, R. B. & Keele, S. W. (1989). Timing functions of the cerebellum. Journal of Cognitive Neuroscience, *1*, 136-152.
- Ivry, R. B., Keele, S. W. & Diener, H. C. (1988). Dissociation of the lateral and medial cerebellum in movement timing and movement execution. Experimental Brain Research, *73*, 167-180.
- Kawato, M. (1996). Trajectory formation in arm movements: Minimization principles and procedures. In H. N. Zelaznik (Ed.), Advances in Motor Learning and Control (pp. 225-259). Champaign, IL.: Human Kinetics.
- Keele, S. W. & Hawkins, H. L. (1982). Explorations of individual differences relevant to high level skill. Journal of Motor Behavior, *14*, 3-23.
- Keele, S. W., Pokorny, R. A., Corcos, D. M., & Ivry, R. (1985). Do perception and motor production share common timing mechanisms: A correlational analysis. Acta Psychologica, *60*, 173-191.
- Kugler, P. N., Kelso, J. A. S., & Turvey, M. T. (1982). On the control and co-ordination of naturally developing systems. In J. A. S. Kelso & J. E. Clark (Eds.), The development of movement control and co-ordination (pp. 5-78). New York: John Wiley & Sons, Ltd.
- Martin, T. A., Keating, J. G., Goodkin, H. P., Bastian, A. J., & Thach, W. T. (1996). Throwing while looking through prisms: I. Focal olivocerebellar lesions impair adaptation. Brain, *119*, 1183-1198.
- Ornstein, R. E. (1969). On the experience of time. Middlesex, England: Penguin Books.
- Peters, M. (1989). The relationship between variability of intertap intervals and interval duration. Psychological Research, *51*, 38-42.
- Robertson, S. D., Zelaznik, H. N., Lantero, D., Bojczyk, K. G., Spencer, R. M., Doffin, J.G., & Schneidt, T. (1999). Correlations for timing consistency among tapping and drawing tasks: Evidence against a single timing process for motor control. Journal of Experimental Psychology: Human Perception and Performance, *25*, 1316-1330.
- Rosenbaum, D. A. (1998). Broadcast theory of timing. In D. A. Rosenbaum & C. E. Collyer (Eds.), Timing of Behavior: Neural, Psychological and Computational Perspective (pp. 215-235). Cambridge, MA.: MIT Press.
- Semjen, A. (1992). Determinants of timing in serial movements. In F. Macar (Ed.), Time, action and cognition: Towards bridging the gap, (pp. 247-261). Dordrecht, Netherlands: Kluwer Academic Publishers.
- Thomas, J. R., & Nelson, J. K. (1996). Research methods in physical activity. Champaign, IL: Human Kinetics.
- Timmann, D., Watts, S., & Hore, J. (1999). Failure of cerebellar patients to time finger opening precisely causes ball high-low inaccuracy in overarm throws. Journal of Neurophysiology, *82*, 103-114.
- Treisman, M., Faulkner, A., & Naish, P. L. (1992). On the relation between time perception and the timing of motor action: Evidence for a temporal oscillator controlling the timing of movement. Quarterly Journal of Experimental Psychology. A, Human Experimental Psychology, *2*, 235-263.
- Turvey, M. T. (1977). Preliminaries to a theory of action with reference to vision. In R. E. S. J. Bransford (Ed.), Perceiving, acting and knowing (pp. 211-265). Hillsdale, NJ: Lawrence Erlbaum.
- Viviani, P., & Terzuolo, C. (1980). Space-time invariance in learned motor skills. In G. E. S. J. Requin (Ed.), Tutorials in motor behavior (pp. 525-533). Amsterdam: North-Holland Publishing Company.
- Viviani, P., & Terzuolo, C. (1982). Trajectory determines movement dynamics. Neuroscience, *7*, 431-437.
- Vorberg, D., & Wing, A. (1996). Modeling variability and dependence in timing. In H. Heuer & S. W. Keele (Eds.), Handbook of perception and action Volume 2: Motor skills (pp. 181-262). New York: Academic Press.
- Wing, A. M. (1980). Timing of movement phases of a repeated response. Journal of Motor Behavior, *12*, 113-124.
- Wing, A.M. (in press). Voluntary timing and brain function: an information processing approach. Brain and Cognition.
- Wing, A., & Kristofferson, A. (1973). Response delays and the timing of discrete motor responses. Perception & Psychophysics, *14*, 5-12.
- Zeiler, M. D. D. (1998). On sundials, springs, and atoms. Behavioural Processes, *44*, 89-99.
- Zelaznik, H. N. & Lantero, D. (1996). The role vision in repetitive circle drawing. Acta Psychologica, *92*, 105-118.
- Zelaznik, H. N., Spencer, R. M., & Doffin, J. (2000). Temporal precision in tapping and circle drawing movements at preferred rates is not correlated: Further evidence against timing as a general purpose ability. Journal of Motor Behavior, *32*, 193-199.

Circle drawing schematic for Experiment 1



Circle drawing schematic for Experiments 2, 3 and 4




Figure 1. A schematic representation of a top view of the circle drawing tasks for the four experiments. Please note that the drawings are not to scale, but are presented to give the reader an understanding of the nature of the circle task, in particular to show the temporal synchronization location.




Figure 2. A representative 4-s slice of behavior in the tapping and intermittent drawing task during the continuation portion of the trial in Experiment 1. These data are not heavily filtered and as such the high frequency noise apparent in the tapping and intermittent drawing trajectory during the pause portion of the movement is evident. The small circles superimposed on the trajectory represents the point in time that is the beginning of cycle n and the end of cycle $n-1$. To the right of the circle trajectory in the y dimension we present the spatial path for the intermittent drawing task for these four seconds.




Figure 3. A representative 4-s slice of behavior in the tapping, continuous drawing and intermittent drawing task during the continuation portion of a trial in Experiment 2. The small circles superimposed on the trajectory represents the point in time that is the beginning of cycle n and the end of cycle $n-1$. The right hand side of panels depict the spatial path of each type of circle drawing movement. The bottom panel shows a four second slice of tapping performance during the continuation portion of a typical trial in Experiment 3. As can be seen from this panel even a 400-ms tapping task has a discernible pause in its movement.